arXiv:1203.6691v1 [math.FA] 30 Mar 2012

APPROXIMATELY MULTIPLICATIVE MAPS FROM WEIGHTED SEMILATTICE ALGEBRAS

YEMON CHOI

28th March 2012

Abstract

We investigate which weighted convolution algebras $\ell^1_\omega(S)$, where S is a semilattice, are AMNM in the sense of Johnson (JLMS, 1986). We give an explicit example where this is not the case. We show that the unweighted examples are all AMNM, as are all $\ell^1_\omega(S)$ where S has either finite width or finite height. Some of these finite-width examples are isomorphic to function algebras studied by Feinstein (IJMMS, 1999).

We also investigate when $(\ell^1_\omega(S), \mathbb{M}_2)$ is an AMNM pair in the sense of Johnson (JLMS, 1988), where $\mathbb{M}_2$ denotes the algebra of 2-by-2 complex matrices. In particular, we obtain the following two contrasting results: (i) for any non-trivial weight on the totally ordered semilattice $\mathbb{N}_{\min}$, the pair $(\ell^1_\omega(\mathbb{N}_{\min}), \mathbb{M}_2)$ is not AMNM; (ii) for any semilattice S, the pair $(\ell^1(S), \mathbb{M}_2)$ is AMNM. The latter result requires a detailed analysis of approximately commuting, approximately idempotent 2×2 matrices.

MSC 2010: 39B72 (primary) 46J10 (secondary)

1 INTRODUCTION

1.1 SETTING THE SCENE

Given a constant $\delta > 0$, we say that a functional ψ on a Banach algebra A is δ-multiplicative if the bilinear map $(a, b) \mapsto \psi(a)\psi(b) - \psi(ab)$ has norm at most δ. It is convenient, thinking of δ as small, to call such functionals approximately multiplicative or almost multiplicative (we shall use the former phrase). Approximately multiplicative functionals have been studied by several authors: an obvious way to obtain examples is to take a multiplicative functional and add a functional of small norm, thought of as a perturbation. The question naturally arises as to whether *all* approximately multiplicative functionals occur in this way.

In [9], B. E. Johnson undertook a systematic study of this phenomenon, and coined the acronym AMNM (for Approximately Multiplicative implies Near Multiplicative). The precise definition will be deferred to a later section. Many examples of commutative Banach algebras with the AMNM property are given in [9], as are some basic hereditary properties. See also [8, 12] for results on uniform algebras, and [6] for results on certain non-uniform function algebras, including $C^k[0,1]^m$. The paper [10] widens the scope of the problem, by considering not just functionals, but approximately multiplicative linear maps between given Banach algebras. This leads to the notion of an AMNM pair; again, the precise definition wil be given below. We note that, as a special case of [10, Theorem 3.1], every amenable Banach algebra has the AMNM property; however, several of the examples in [6, 8, 9] possess non-zero point derivations, so that amenability is far from necessary for AMNM.

In this paper, we investigate these AMNM problems for the weighted ℓ^1-convolution algebras of semilattices. Such algebras have provided provide useful test cases for various conjectures and techniques concerning commutative Banach algebras. Moreover, any weighted semilattice algebra contains a dense subalgebra spanned by commuting idempotents. Thus, some of our work can be viewed as continuing an old strand of Banach algebra theory, which considers lifting and perturbation questions for families of idempotents. The main difference here is that we are not limiting ourselves to families of pairwise orthogonal idempotents, but allowing more complicated order structure.

The original motivation for the present work arises from studying the cases where the underlying semilattice is $\mathbb{N}_{\min}$, the set of natural numbers equipped with pairwise minimum as a semigroup product. The weighted ℓ^1-algebras of $\mathbb{N}_{\min}$ turn out to be isomorphic to function algebras that were studied by J. F. Feinstein in [2]; they have also been studied in the context of certain generalized notions of amenability, see for instance [1, §3.10]. Moreover, some of these algebras satisfy such versions of amenability while having non-trivial 2nd-degree simplicial cohomology (the present author, unpublished calculations).

1.2 OVERVIEW OF THE PAPER

We have tried to make this paper self-contained, save for some basic knowledge of Banach algebras. Thus, in Section 2 we give the relevant definitions of the AMNM property for algebras and for pairs of algebras, as promised earlier; and we record some basic observations on convolution algebras and their characters. We then observe that $\ell^1(S)$ is AMNM for any semilattice S (Theorem 3.1). On the other hand, we give an explicit example of a semilattice T and a weight on T such that the weighted convolution algebra $\ell^1_\omega(T)$ is not AMNM (Theorem 3.4). In Section 3.3, as a special case of a general technical result, we prove that if S has either finite width or finite height, then $\ell^1_\omega(S)$ is AMNM for every weight ω. This applies in particular when $S = \mathbb{N}_{\min}$, the original case of interest.

The picture is far less complete if we consider approximately multiplicative maps into algebras other than $\mathbb{C}$. In Theorem 4.2, we show that if A is the convolution algebra for any non-trivial weight on $\mathbb{N}_{\min}$, then the pairs $(A, \mathbf{T}_2)$ and $(A, \mathbb{M}_2)$ are not AMNM, where $\mathbb{M}_2$ is the algebra of 2×2 matrices and $\mathbf{T}_2$ is the (commutative, non-semisimple) algebra of dual numbers over $\mathbb{C}$.

These examples suggest that, if we want positive AMNM results for range algebras other than $\mathbb{C}$, we should focus attention on the unweighted case. Indeed, we prove that for an arbitrary semilattice S, the pairs $(\ell^1(S), \mathbf{T}_2)$ and $(\ell^1(S), \mathbb{M}_2)$ are "uniformly AMNM" (the terminology is explained below, in Definition 2.4). The proof of this for $\mathbb{M}_2$ takes up all of Section 5: although the techniques used are elementary, a complete proof seems to require substantially more work than is needed for $\mathbf{T}_2$. Finally, we close the paper by briefly discussing some possible avenues for future work.

Remark 1.1. Some of our calculations would work for certain weighted algebras on abelian Clifford semigroups, specifically those where the weight is trivial on each group component. We have decided to only consider the semilattice case for now: an adequate treatment of weighted abelian Clifford semigroups would require a more detailed look at AMNM problems for Beurling algebras than the present paper can accommodate. The same remarks apply for inverse semigroups: we did not see how to go beyond superficial generalizations of the results here.

Remark 1.2. After completing the work presented in this article, we learned of the paper [11], which considers a related but different notion of AMNM (briefly, the order of quantifiers is different). Although the stability result that is proved in [Theorems 2 and 5, ibid.] is different from ours, it may be possible to use those arguments to streamline the approach taken in Section 5, and perhaps even to extend the results of Section 5 from $\mathbb{M}_2$ to $\mathbb{M}_n$.

2 PRELIMINARIES

We assume familiarity with the basics of Banach spaces and Banach algebras. Throughout $\widehat{\otimes}$ denotes the projective tensor product of Banach spaces. If A is a Banach algebra then $\pi_A : A \widehat{\otimes} A \to A$ denotes the unique bounded bilinear map satisfying $\pi_A(a_1 \otimes a_2) = a_1 a_2$.

2.1 DEFINING AMNM

Our notation is different from that of Johnson's articles [9, 10], and so we repeat some of the basic definitions for sake of clarity.

Definition 2.1 (Multiplicative defect)**.** Let A and B be Banach algebras, and let $T : A \to B$ be a bounded linear map. We define the multiplicative defect of T to be

$$\begin{aligned} \operatorname{def}(T) &= \| T \circ \pi_A - \pi_B \circ (T \otimes T) : A \widehat{\otimes} A \to B \| \\ &= \sup\{ \| T(xy) - T(x)T(y) \| \ : \ x, y \in A, \|x\| \leq 1, \|y\| \leq 1 \}. \end{aligned} \tag{2.1}$$

T is said to be δ-multiplicative, in the sense of [10], when $\operatorname{def}(T) \leq \delta$. Note that Johnson uses the notation $T^{\vee}$ instead of $\operatorname{def}(T)$. Of course, a 0-multiplicative map is just one that is multiplicative in the usual sense, that is, a continuous linear algebra homomorphism (which need not be unital, even if A and B are unital algebras). We denote by $\operatorname{Mult}(A, B)$ the set of multiplicative, bounded linear maps $A \to B$; a non-zero element of $\operatorname{Mult}(A, \mathbb{C})$ is called a character of A.

Given a subset K in a metric space (X, d) and $y \in X$, define $\operatorname{dist}_X(y, K)$ to be $\inf_{x \in X} d(x, y)$.

Definition 2.2 (AMNM algebras, [9])**.** A Banach algebra A is said to be AMNM, or have the AMNM property, or have stable characters, if for each $\varepsilon > 0$ there exists $\delta > 0$ such that

$$\psi \in A^*, \ \operatorname{def}(\psi) \leq \delta \implies \operatorname{dist}_{A^*}(\psi, \operatorname{Mult}(A, \mathbb{C})) \leq \varepsilon.$$

While Definition 2.2 does not require A to be commutative, it seems most natural to study the AMNM property for commutative Banach algebras, since those are the ones for whom characters are most informative (via Gelfand theory). For non-commutative algebras A, the following definition seems more natural.

Definition 2.3 (AMNM pairs of Banach algebras, [10])**.** Let A and B be Banach algebras. We say that the pair (A, B) is an AMNM pair if, for every $K > 0$ and $\varepsilon > 0$, there exists $\delta > 0$ such that

$$T \in \mathcal{L}(A, B), \ \|T\| \leq K, \ \operatorname{def}(T) \leq \delta \implies \operatorname{dist}_{\mathcal{L}(A,B)}(T, \operatorname{Mult}(A, B)) \leq \varepsilon.$$

The presence of the *a priori* upper bound K may seem curious at first sight. One reason for imposing such a bound is that otherwise we can find T such that $\operatorname{dist}_{\mathcal{L}(A,B)}(T, \operatorname{Mult}(A,B))$ is arbitrarily small while $\operatorname{def}(T) \geq 1$. See [10, p. 295] for an example with $A = \mathbb{C}$ and B the algebra $\mathbb{M}_2$ of 2×2 matrices.

Nevertheless, in the present paper, we shall obtain examples of pairs of algebras which are not just AMNM, but which satisfy a stronger property, defined as follows.

Definition 2.4. Let A and B be Banach algebras. We say that (A,B) is a uniformly AMNM pair if, for every $\varepsilon > 0$, there exists $\delta > 0$ such that

$$T \in \mathcal{L}(A,B),\ \operatorname{def}(T) \leq \delta \implies \operatorname{dist}_{\mathcal{L}(A,B)}(T, \operatorname{Mult}(A,B)) \leq \varepsilon.$$

With this terminology, a Banach algebra A is AMNM if and only if the pair $(A, \mathbb{C})$ is uniformly AMNM. In some cases, AMNM pairs are automatically uniformly AMNM: for instance, this is the case when $B = C(X)$, since a δ-multiplicative linear map $A \to C(X)$ has norm at most $1 + \delta$ [7, Proposition 5.5]. Note that Johnson [10, Example 1.5] has given an example of a commutative, semisimple Banach algebra B for which the pair $(\mathbb{C}, B)$ is AMNM but not uniformly AMNM.

2.2 SOME NOTATION AND TERMINOLOGY

A weight on a set S is a function $\omega : S \to (0,\infty)$. Given such a weight ω, we write $\ell^1_\omega(S)$ for the corresponding weighted ℓ^1-space. Throughout this paper, whenever we refer to a weight on a semigroup, we always mean a *submultiplicative* weight. By a weighted semigroup, we mean a pair (S,ω) where S is a semigroup and ω is a weight on S. A routine calculation shows that if (S,ω) is a weighted semigroup, then $\ell^1_\omega(S)$, equipped with the natural convolution product, is a Banach algebra, called the weighted semigroup algebra or convolution algebra of (S,ω).

A bounded function f from a set S to a Banach space B has a unique continuous extension to a bounded linear map $\ell^1(S) \to B$, whose norm is precisely $\sup_{t\in S} \|f(t)\|$. Moreover, if S is a semigroup and B is a Banach algebra, this extension will be multiplicative if and only if the original function f is multiplicative. Thus, the characters of $\ell^1(S)$ are in natural bijection with the non-zero semigroup homomorphisms $S \to (\mathbb{C}, \times)$, sometimes called the semi-characters of S.

The corresponding version for weighted semigroup algebras is equally straightforward: bounded linear maps $\ell^1_\omega(S) \to B$ correspond to functions $f : S \to B$ such that

$$\sup_{s\in S} \omega(s)^{-1}\|f(s)\| < \infty, \tag{2.2}$$

and so forth. Such a function will be called an ω-bounded map from S to B. In the case where $B = \mathbb{C}$ and $f : S \to \mathbb{C}$, the quantity in (2.2) will be denoted by $\|f\|_{\infty,\omega^{-1}}$.

Since we can naturally and isometrically identify $\ell^1_\omega(S) \widehat{\otimes} \ell^1_\omega(S)$ with $\ell^1_{\omega\times\omega}(S \times S)$, it is easy to check that for a given function $f : S \to B$, the multiplicative defect of the corresponding linear map $\ell^1_\omega(S) \to B$ is

$$\operatorname{def}_\omega(f) := \sup_{x,y\in S} \frac{\|f(x)f(y) - f(xy)\|}{\omega(x)\omega(y)}.$$

In this way, all AMNM questions where the domain algebra A is a weighted semigroup algebra can be rephrased in terms of ω-bounded maps from the given semigroup and $(\omega \times \omega)$-bounded maps from its Cartesian square. This is sometimes convenient if we need to define a map via case-by-case checking.

2.3 WEIGHTED SEMILATTICE ALGEBRAS, AND THEIR CHARACTERS

A semilattice is a commutative semigroup in which each element is idempotent. Even quite simple semilattices can give rise to interesting Banach algebras, once we allow for weights. The following examples provided the initial motivation for this article, and will be revisited in Theorem 4.2.

Example 2.5 ("Feinstein algebras")**.** Let $\mathbb{N}_{\min}$ denote the semilattice obtained by equipping the set of natural numbers with the binary operation $(m,n) \mapsto \min(m,n)$. The weights on this semilattice are precisely the functions $\omega : \mathbb{N} \to [1,\infty)$. The convolution algebra $\ell^1_\omega(\mathbb{N}_{\min})$ is semisimple and its character space can be naturally identified with $\mathbb{N}$. (This is alluded to, without details, in [4, 11.1.5]; see also Lemma 2.8 below.) In fact, the Gelfand transform maps $\ell^1_\omega(\mathbb{N}_{\min})$ onto a dense subalgebra $B_\omega \subset c_0(\mathbb{N})$, defined by

$$B_\omega = \{f \in c_0(\mathbb{N}) \ : \ |f_1|\omega(1) + \sum_{j\geq 2} |f_{j+1} - f_j|\omega(j) < \infty\}. \tag{2.3}$$

equipped with the obvious norm. The unitizations of the algebras B_ω are isomorphic to examples studied in [2]. Note that in the cases where ω is bounded, $\ell^1_\omega(\mathbb{N}_{\min})$ is isomorphic as a Banach algebra to $\ell^1(\mathbb{N}_{\min})$, and B_ω will consist of those c_0-sequences which have bounded variation. For more details, see [1, §3].

Remark 2.6. In the literature, terminology and notation for these examples has varied. Strictly speaking, Feinstein's construction in [2] is more general: he defines, for any given sequence $\alpha = (\alpha_n)_{n\geq 1}$ of strictly positive real numbers, a commutative unital Banach algebra $A_\alpha \subset C(\mathbb{N} \cup \{\infty\})$, and in his notation our B_α coincides with $A_\alpha \cap c_0$. Note, however, that in some later papers the algebra A_α is defined to be what we have denoted by B_α.

For general semilattices, a systematic approach to the character space is via the language of filters. First we introduce some notation that will be used later in Section 3.3. Given a semilattice S and a subset $E \subseteq S$, we define the following sets:

$$\langle E \rangle_n = \{x_1 \cdots x_n \ : \ x_1, \ldots, x_n \in E\} \quad (n \geq 1), \tag{2.4}$$

and $\langle E \rangle = \bigcup_{n\geq 1} \langle E \rangle_n$. Note that $\langle E \rangle$ is the sub-semilattice of S generated by E.

Definition 2.7. Let S be a semilattice and let $F \subseteq S$. We say that F is a filter in S if it satisfies the following three properties: it is non-empty; it is closed under multiplication (i.e. $xy \in F$ whenever $x, y \in F$); and it is upwards closed in S (i.e. whenever $x \in S$ and $y \in F$ with $xy \in F$, then $x \in F$).

Given a non-empty subset E in a semilattice S, there exists a smallest filter in S which contains E. This can be described concretely: it is the set

$$F = \bigcup_{y \in \langle E \rangle} \{x \in S \ : \ x \succeq y\}. \tag{2.5}$$

The following lemma assembles some basic results, which are well known, at least implicitly, for multiplicative linear functions on semilattice algebras. (See e.g. [4, 11.1.1].) Since the proofs are no harder for the weighted versions, we leave them to the reader.

Lemma 2.8. *Let S be a semilattice and $F \subset S$. The following are equivalent:*

(i) *F is a filter in S;*

(ii) *the indicator function of F is a character on $\ell^1_\omega(S)$ for every submultiplicative weight ω;*

(iii) *there exist a submultiplicative weight ω such that the indicator function of F is a character on $\ell^1_\omega(S)$.*

Moreover, if $\phi : S \to \mathbb{C}$ is multiplicative and non-zero, and $\phi = 1$ on some subset $E \subseteq S$, then $\phi = 1$ on any filter containing E.

3 THE AMNM PROPERTY FOR WEIGHTED SEMILATTICE ALGEBRAS

3.1 THE UNWEIGHTED CASE, AS A GUIDE

Theorem 3.1. *Let S be a semilattice. Then $\ell^1(S)$ is AMNM.*

This result, or equivalent reformulations, may have been implicitly known to previous authors. We shall give a complete proof, since it allows us to introduce some basic results and techniques which will be instructive for later arguments.

Define $f : [0, 1/4) \to [0, 1/2)$ by

$$f(t) = \frac{1}{2}\left(1 - \sqrt{1-4t}\right). \tag{3.1}$$

The following properties are straightforward to verify.

- f is convex and monotone increasing, and $f(0) = 0$.
- $\rho : t \mapsto t^{-1} f(t)$ is monotone increasing on $(0, 1)$, and $\lim_{t\searrow 0} t^{-1} f(t) = 1$.
- $f(2/9) = 1/3$ and $\rho(2/9) = 3/2$.

Notation. Given $r \geq 0$ and $w \in \mathbb{C}$, denote the closed disc of radius r and centre w by $\overline{\mathbb{D}}_w(r)$. If $r > 0$, denote the open disc of radius r and centre w by $\mathbb{D}_w(r)$.

Lemma 3.2. *Let $z \in \mathbb{C}$, let $\varepsilon \in [0, 2/9]$, and suppose $z^2 - z \in \overline{\mathbb{D}}_0(\varepsilon)$. Then $\operatorname{dist}_{\mathbb{C}}(z, \{0,1\}) \leq f(\varepsilon) \leq \frac{3}{2}\varepsilon$.*

Proof. Put $w = z - z^2$. Then $\left(z - \frac{1}{2}\right)^2 = \frac{1}{4}(1 - 4w)$, and so $z - \frac{1}{2} = \pm\frac{1}{2}\sqrt{1-4w}$ (taking the branch of the square root function for which $z \to 0$ as $w \to 0$). It follows that

$$\operatorname{dist}_{\mathbb{C}}(z, \{0,1\}) \leq \left| \frac{1}{2} - \frac{1}{2}\sqrt{1-4w} \right|.$$

The Taylor expansion (about 0) of the function $w \mapsto 1 - \sqrt{1-4w}$ has non-negative coefficients. Hence $\operatorname{dist}_{\mathbb{C}}(z, \{0,1\}) \leq f(|w|)$, and the rest follows from our earlier observations. $\square$

Theorem 3.1 follows immediately from the following technical result.

Proposition 3.3. *Let S be a semilattice, and let $\psi : S \to \mathbb{C}$ satisfy* $\operatorname{def}(\psi) < 2/9$. *Let*

$$S_1 = \psi^{-1}\left(\mathbb{D}_1(1/3)\right) = \{e \in S \ : \ |\psi(e) - 1| < 1/3\},$$

and let χ be the indicator function of S_1. Then χ is a multiplicative function $S \to \mathbb{C}$, satisfying

$$\|\psi - \chi\|_\infty \leq 3\|\psi^2 - \psi\|_\infty \leq \frac{3\delta}{2}. \tag{3.2}$$

Proof. By Lemma 3.2, we have

$$\sup_{e\in S} \min(|\psi(e) - 1|, |\psi((e)|) \leq \sup_{e\in S} \frac{3}{2}|\psi(e)^2 - \psi(e)| = \frac{3}{2}\operatorname{def}(\psi) < \frac{1}{3}.$$

Hence, if we set $S_0 = \psi^{-1}(\mathbb{D}_0(1/3))$, then $S = S_0 \sqcup S_1$, from which the inequality (3.2) follows immediately.

To prove χ is multiplicative, it suffices by Lemma 2.8 to show that S_1 is either empty or a filter. Suppose that S_1 is non-empty. If $e, f \in S_1$, then

$$|\psi(e)\psi(f)| > \left(\frac{2}{3}\right)^2 = \frac{4}{9}.$$

Hence $|\psi(ef)| > 4/9 - \operatorname{def}(\psi) > 1/3$, forcing $ef \in S \setminus S_0 = S_1$. If $e \succeq f \in S$ and $f \in S_1$, then since $|\psi(f)|^{-1} < 3/2$, we have

$$|\psi(e) - 1| \leq \frac{3}{2}|\psi(e)\psi(f) - \psi(f)| = \frac{3}{2}\operatorname{def}(\psi) < \frac{1}{3},$$

forcing $e \in S_1$. Thus S_1 is a filter, and the proof is complete. ☐

3.2 A WEIGHTED SEMILATTICE WHICH IS NOT AMNM

Theorem 3.4. *There exists a locally finite semilattice T and a submultiplicative weight ω on T such that $\ell^1_\omega(T)$ is not AMNM.*

The counter-example T will be built out of copies of free semilattices. Given a set S, let $\mathbf{2}_*^{[S]}$ denote the free semilattice generated by S; this can be identified with the set of all non-empty finite subsets of S, with semigroup product given by union. For our purposes it is more convenient to regard elements of $\mathbf{2}_*^{[S]}$ as reduced words in the generators. There is a natural length function $\gamma_S : \mathbf{2}_*^{[S]} \to \mathbb{N}$, where $\gamma_S(x)$ is the minimum number of elements in S needed to generate x. A little thought shows that $\gamma_S(xy) \leq \min(|S|, \gamma_S(x) + \gamma_S(y))$ for all $x, y \in \mathbf{2}_*^{[S]}$.

If S is a finite set, then $\mathbf{2}_*^{[S]}$ has a zero (i.e. minimum) element, namely the product of all elements of S; we denote this element by θ_S. Note that $\gamma_S(\theta_S) = |S|$.

Proposition 3.5. *Let S be a finite set of cardinality ≥ 2. Fix a constant $C > 1$, and define $\omega_S : \mathbf{2}_*^{[S]} \to [1, \infty)$ by*

$$\omega_S(e) = \begin{cases} C^{\gamma_S(e)} & \text{if } e \neq \theta_S \\ C & \text{if } e = \theta_S \end{cases}$$

Then ω_S is a submultiplicative weight on $\mathbf{2}_*^{[S]}$. Moreover, if we define $\psi : \mathbf{2}_*^{[S]} \to \{0,1\}$ by

$$\psi(e) = \begin{cases} 1 & \text{if } e \neq \theta_S, \\ 0 & \text{if } e = \theta_S, \end{cases}$$

then:

(i) *as an element of $\ell^1_{\omega_S}(\mathbf{2}_*^{[S]})^*$, $\operatorname{def}(\psi) \leq C^{-|S|}$;*

(ii) *if $\phi : \mathbf{2}_*^{[S]} \to \{0,1\}$ is multiplicative, then*

$$\sup_{e \in \mathbf{2}_*^{[S]}} \omega_S(e)^{-1} |\psi(e) - \phi(e)| \geq C^{-1}.$$

Proof. Since γ_S is subadditive and $\omega_S(e) \leq C^{\gamma_S(e)}$ for all $e \in \mathbf{2}_*^{[S]}$, it is clear that ω_S is submultiplicative.

To prove (i), let $e, f \in \mathbf{2}_*^{[S]}$. If $ef \neq \theta_S$ then $\psi(e)\psi(f) = 1 = \psi(ef)$. If $e = \theta_S$ or $f = \theta_S$ then $\psi(e)\psi(f) = 0 = \psi(ef)$. The only remaining cases are those where $e, f \neq \theta_S$ while $ef = \theta_S$. In such cases, we have $\gamma_S(e) + \gamma_S(f) \geq |S|$, and so

$$|\psi(e)\psi(f) - \psi(ef)| = 1 \leq \delta C^{-|S|} \omega(e)\omega(f).$$

Thus in all cases, $|\psi(e)\psi(f) - \psi(ef)| \leq C^{-|S|}\omega(e)\omega(f)$.

To prove (ii), let $\phi : S \to \{0,1\}$ be multiplicative. If $\phi(\theta_S) = 1$ then

$$\omega(\theta_S)^{-1} |\phi(\theta_S) - \psi(\theta_S)| = C^{-1}.$$

If not, then $\phi(\theta_S) = 0$. Therefore, since ϕ is multiplicative and S generates $\mathbf{2}_*^{[S]}$, there exists $s_0 \in S$ with $\phi(s_0) = 0$, and so

$$\omega(s_0)^{-1} |\phi(s_0) - \psi(s_0)| = C^{-1}.$$

□

If $(F_i)_{i \in \mathbb{I}}$ is a family of semilattices, consider the set $\{\theta\} \sqcup \coprod_{i \in \mathbb{I}} F_i$, where θ is a formal symbol. This can be made into a semilattice if we define the product as follows: θ is an absorbing zero element; the product of two elements in F_i is their usual product; the product of elements in F_i and F_j is θ whenever $i \neq j$. We call this semilattice the orthogonal direct sum of the family $(F_i)_{i \in \mathbb{I}}$.

Proof of Theorem 3.4. Let $(F_n)_{n\geq 1}$ be a sequence of finite sets with $|F_n| \nearrow \infty$. Define T to be the orthogonal direct sum of the family $(\mathbf{2}_*^{[F_n]})_{n\in\mathbb{N}}$, with θ being the zero element of T. To ease notation, denote the zero element of $\mathbf{2}_*^{[F_n]}$ by θ_n, and denote the length function of $\mathbf{2}_*^{[F_n]}$ by γ_n.

Fix a constant $C > 1$, and define $\omega : T \to [1,\infty)$ by

$$\omega(\theta) := 1 \quad , \quad \omega(\theta_n) := C \text{ for each } n \in \mathbb{N} \quad , \quad \omega(e) := C^{\gamma_n(e)} \quad \text{if } e \in \mathbf{2}_*^{[F_n]} \setminus \{\theta_n\}.$$

Define $\psi_n : T \to \{0,1\}$ by

$$\psi_n(e) = \begin{cases} 1 & \text{if } E \in S_n \setminus \{\theta_n\}, \\ 0 & \text{otherwise.} \end{cases}$$

If $\phi : T \to \{0,1\}$ is multiplicative, then by part (ii) of Proposition 3.5,

$$\|\psi_n - \phi\| \geq \sup\{\omega(e)^{-1}|\psi_n(e) - \phi(e)| \ : \ e \in \mathbf{2}_*^{[F_n]}\} \geq C^{-1}.$$

Hence,

$$\operatorname{dist}_{\ell^1_\omega(T)^*}(\psi_n, \operatorname{Mult}(\ell^1_\omega(T), \mathbb{C})) \geq C^{-1} \quad \text{for all } n.$$

On the other hand, by Proposition 3.5(i),

$$\operatorname{def}_\omega(\psi_n) = \sup\left\{ \frac{|\psi_n(e)\psi_n(f) - \psi_n(ef)|}{\omega(e)\omega(f)} \ : \ e, f \in \mathbf{2}_*^{[F_n]} \right\} \leq C^{-|F_n|} \to 0.$$

Thus $\ell^1_\omega(T)$ is not AMNM. ☐

The example (T, ω) was found while trying to prove that all weighted semilattice algebras are AMNM, and realizing that the attempted proof only worked when one could verify a certain technical condition on a given weighted semilattice (S, ω). This condition, and the proof that it suffices to ensure $\ell^1_\omega(S)$ is AMNM, will be our next topic.

3.3 WEIGHTED SEMILATTICES WHICH ARE AMNM

The following lemma is a substitute for Lemma 3.2. It is less informative in the case of $\ell^1(S)$, but is more convenient in weighted cases.

Lemma 3.6. *Fix $\delta > 0$ and $c > 1$. Let A be a Banach algebra, e an idempotent in A, and $\psi \in A^*$. Then*

$$\min(|\psi(e)|, |1 - \psi(e)|) \leq \operatorname{def}(\psi)^{1/2}\|e\|. \tag{3.3}$$

Proof. If $e = 0$ this is trivial. If $e \neq 0$, observe that

$$\operatorname{def}(\psi) \geq \frac{|\psi(e) - \psi(e)^2|}{\|e\|^2} = \frac{|\psi(e)|}{\|e\|}\frac{|1 - \psi(e)|}{\|e\|} \geq \left(\frac{\min(|\psi(e)|, |1 - \psi(e)|)}{\|e\|}\right)^2,$$

and (3.3) follows. ☐

By (3.3) and the definition of the norm on $\ell^1_\omega(S)$, we see that if $\psi : S \to \mathbb{C}$ is ω-bounded with $\operatorname{def}_\omega(\psi) \leq \delta$, then there exists an ω-bounded function $\phi : S \to \{0,1\}$ such that $\|\psi - \phi\|_{\infty,\omega^{-1}} \leq \delta^{1/2}$. Also, since

$$|\phi(s)\phi(t) - \phi(st)| \leq \begin{cases} |\phi(s)||\phi(t) - \psi(t)| + |\phi(s) - \psi(s)||\psi(t)| \\ +|\psi(s)\psi(t) - \psi(st)| + |\psi(st) - \phi(st)|\,, \end{cases}$$

we have $\operatorname{def}_\omega(\phi) \leq 3\delta^{1/2} + \delta$. From this, routine arguments (which we omit) yield the following necessary and sufficient condition for $\ell^1_\omega(S)$ to be AMNM.

Corollary 3.7. *Let (S,ω) be a weighted semilattice. The following are equivalent:*

(i) *$\ell^1_\omega(S)$ is AMNM;*

(ii) *For any $\varepsilon > 0$, there exists $\delta > 0$, such that whenever $\phi : S \to \{0,1\}$ satisfies*

$$\sup_{e,f\in S} \frac{|\phi(ef) - \phi(e)\phi(f)|}{\omega(e)\omega(f)} \leq \delta\,,$$

then there exists a subset $F \subseteq S$, either empty or a filter, such that

$$\sup_{e\in S} \frac{|\chi_F(e) - \phi(e)|}{\omega(e)} \leq \varepsilon.$$

The following terminology is non-standard, since we have not been able to find prior terminology in the literature.

Definition 3.8 (Verbosity and prolixity)**.** Let E be a subset of a semilattice S. The verbosity of E, denoted $\operatorname{vb}(E)$, is defined to be $\inf\{n \in \mathbb{N} : \langle E\rangle_N = \langle E\rangle\}$, with the usual convention that the infimum of the empty set is $+\infty$. We define the prolixity of S to be $\sup_{E\subseteq S} \operatorname{vb}(E)$, and denote it by $\operatorname{prol}(S)$.

Here, the sets $\langle E\rangle_n$ and $\langle E\rangle$ are as defined in (2.4). Note that the condition $\operatorname{prol}(S) < \infty$ does not imply S is finitely generated as a semigroup. (Indeed, finitely generated semilattices are finite.)

Example 3.9. Let S be the free semilattice on n generators. Then $\operatorname{prol}(S) = n$.

Example 3.10. If S has width $\leq n$ as a partially ordered set (i.e. there exist n chains in S whose union is all of S) then $\operatorname{prol}(S) \leq n$.
(Fix chains $C_1, \dots, C_n$ in S such that $S = \bigcup_{i=1}^n C_i$. Let $E \subseteq S$ and $x \in \langle E\rangle$. For some integer k there exists $y_1, \dots, y_k \in E$ such that $x = y_1 \cdots y_k$. Partition $\{1, \dots, k\}$ into disjoint, non-empty subsets $J(1), \dots, J(m)$, where $m \leq n$, such that $y_r \in C_i$ for all $r \in J(i)$. For each i, the set $\{y_r : r \in J(i)\}$ is totally ordered (as a subset of S) and so has a least element, say $y_{s(i)}$. Then $\prod_{r\in J(i)} y_r = y_{s(i)}$, so

$$x = y_{s(1)} \dots y_{s(m)} \in \langle E\rangle_m \subseteq \langle E\rangle_n$$

and thus $\operatorname{vb}(E) \leq n$.)

Example 3.11. Let $n \geq 2$. If S has height $\leq n$ (i.e. each chain in S has cardinality $\leq n$) then $\operatorname{prol}(S) \leq n$.
(Let $E \subseteq S$ and $x \in \langle E \rangle$. For some integer k there exists $y_1, \ldots, y_k \in E$ such that $x = y_1 \cdots y_k$. Clearly $y_1 \succeq y_1 y_2 \succeq \cdots \succeq y_1 y_2 \cdots y_k$. Let

$$J = \{r \in \{2, \ldots, k\} \ : \ y_1 \cdots y_{r-1} \neq y_1 \cdots y_r\},$$

and enumerate its elements in increasing order as $j_1 < \ldots j_m$. Since chains in S have cardinality at most n, we have $m \leq n+1$, and thus $\operatorname{vb}(E) \leq n$.)

We need one more technical definition. Let (S, ω) be a weighted semilattice, and let C, α be constants ≥ 1. We say that the pair (S, ω) is (C, α)-governed if

$$\max_{y \in \langle E \rangle} \omega(y) \leq C \max_{x \in E} \omega(x)^\alpha \quad \text{for each subset } E \subseteq S.$$

This condition is somewhat artificial, and is set up to make the proof of the next theorem work. Let us first consider some examples.

Example 3.12.

(i) If $\sup_{x \in S} \omega(x) = C < \infty$, then (S, ω) is $(C, 1)$-governed.

(ii) If $\operatorname{prol}(S) = n < \infty$, then (S, ω) is $(1, n)$-governed for every weight ω.

(iii) Let (T, ω) be the weighted semilattice constructed in the proof of Theorem 3.4. Then there are no constants K and α such that (T, ω) is $(K, \alpha$-governed.

Theorem 3.13. *Let (S, ω) be a weighted semilattice, which is (C, α)-governed for some constants $C \geq 1$ and $\alpha \geq 1$. Then $\ell^1_\omega(S)$ is AMNM.*

As a special case of Theorem 3.13, we get another proof that $\ell^1(S)$ is AMNM for every semilattice S (see Example 3.12(i)). More interestingly, we can deduce that if S is a semilattice with either finite width or finite height, then $\ell^1_\omega(S)$ is AMNM for every submultiplicative weight ω (see Example 3.12(ii)). In particular, for any weight function $\omega : \mathbb{N} \to [1, \infty)$, the algebra B_ω from Example 2.5 is AMNM, as it is isomorphic to $\ell^1_\omega(\mathbb{N}_{\min})$.

By Corollary 3.7, Theorem 3.13 will follow from the following more technical result.

Proposition 3.14. *Let (S, ω) be a weighted semilattice that is (C, α)-governed. Let $\delta \in (0,1)$ and $K > \max(C, 2^{1+1/\alpha})$. Suppose $\psi : S \to \{0,1\}$ satisfies*

$$|\psi(st) - \psi(s)\psi(t)| \leq \delta \omega(s) \omega(t) \qquad \text{for all } s, t \in S.$$

Then there exists $F \subseteq S$, either empty or a filter, such that

$$|\chi_F(x) - \psi(x)| \leq K \delta^{1/(\alpha+1)} \omega(x) \quad \text{for all } x \in S.$$

Proof of Proposition 3.14. Let

$$S_{\text{fix}} := \{x \in S \ : \ \omega(x) \leq (K\delta)^{-1/(\alpha+1)}\}$$

(we think of this as the set where ω is small). Let $E := \{y \in S_{\text{fix}} \ : \ \psi(y) = 1\}$ and let

$$F := \bigcup_{y \in \langle E \rangle} \{x \in S \ : \ x \succeq y\}.$$

Recall: if E is empty then so is F; otherwise, F is the filter generated by E in S.

To motivate the next step, suppose $\phi : S \to \mathbb{C}$ is multiplicative and is close in norm to ψ. Then (provided δ is sufficiently small) we must have $\phi = 1$ on E; and so, as remarked in Lemma 2.8, ϕ must be 1 on F. Therefore, ψ must also be 1 on $F \cap S_{\text{fix}}$, i.e. we must have $E = F \cap S_{\text{fix}}$.

So, our next step is to verify that $E = F \cap S_{\text{fix}}$. It suffices to show that $F \cap S_{\text{fix}} \subseteq E$, and the proof of this goes via the following claim:

CLAIM. Let $k \in \mathbb{N}$. Then $\psi(y) = 1$ for all $y \in \langle E \rangle_k$.

PROOF OF CLAIM. Induction on k. If $k = 1$ there is nothing to prove. Suppose that the claim holds for $k = m - 1$ where $2 \leq m$, and let $x_1, \ldots, x_m \in E$. Put $y' = x_1 \cdots x_{m-1}$; then by the inductive hypothesis, and the assumption that (S, ω) is (C, α)-governed,

$$\begin{aligned}
|\psi(y'x_m) - 1| &= |\psi(y'x_m) - \psi(y')\psi(x_m)| \\
&\leq \delta\omega(y')\omega(x_m) \\
&\leq \delta C \left((K\delta)^{-1/(\alpha+1)} \right)^{\alpha} (K\delta)^{-1/(\alpha+1)} = C/K < 1
\end{aligned}$$

which, since ψ is 0-1-valued, forces $\psi(y'x_m) = 1$, so that the claim holds for $k = m$.

Now let $z \in F \cap S_{\text{fix}}$. Since $z \in F$, there exists $k \in \mathbb{N}$ and $y \in \langle E \rangle_k$ such that $zy = y$. By our claim, $\psi(y) = 1$. Since $z \in S_{\text{fix}}$ this implies

$$\begin{aligned}
|\psi(z) - 1| &= |\psi(y)\psi(z) - \psi(yz)| \\
&\leq \delta\omega(y)\omega(z) \\
&\leq \delta \cdot C(K\delta)^{-1/(\alpha+1)}(K\delta)^{-1/(\alpha+1)} \\
&= CK^{-2/(\alpha+1)}\delta^{(\alpha-1)/(\alpha+1)} \leq C/K < 1
\end{aligned}$$

forcing $\psi(z) = 1$, so that $z \in E$.

Thus $F \cap S_{\text{fix}} = E$, as required. We therefore have $S = E \cup (S_{\text{fix}} \setminus F) \cup (S \setminus S_{\text{fix}})$. Now observe that:

- when $x \in E$ then $|\chi_F(x) - \psi(x)| = 0$;
- when $x \in S_{\text{fix}} \setminus F \subseteq S_{\text{fix}} \setminus E$ then $\chi_F(x) = 0 = \psi(x)$, so that $|\chi_F(x) - \psi(x)| = 0$;
- when $x \in S \setminus S_{\text{fix}}$ then $|\chi_F(x) - \psi(x)| \leq 2 \leq K^{\alpha/(\alpha+1)} < K\delta^{1/(\alpha+1)}\omega(x)$.

Putting these cases together, the proposition is proved. $\square$

4 MORE GENERAL RANGE ALGEBRAS

For which Banach algebras B and weighted semilattices (S, ω) is $(\ell^1_\omega(S), B))$ an AMNM pair? Since we do not have a complete answer in the case $B = \mathbb{C}$, we can expect only partial results in the more general case. A natural place to start is with those semilattices covered by Theorem 3.13, in particular with $\mathbb{N}_{\min}$. Recall that by this theorem, $\ell^1_\omega(\mathbb{N}_{\min})$ is AMNM for any weight ω. In contrast, we will now show that whenever ω is a non-trivial (=unbounded) weight on $\mathbb{N}_{\min}$, the pairs $(\ell^1_\omega(\mathbb{N}_{\min}), \mathbf{T}_2)$ and $(\ell^1_\omega(\mathbb{N}_{\min}), \mathbb{M}_2)$ fail to be AMNM. Here, $\mathbb{M}_2$ is the algebra of 2×2 complex matrices, and $\mathbf{T}_2$ is the subalgebra

$$\left\{ \begin{pmatrix} a & b \\ 0 & a \end{pmatrix} \ : \ a, b \in \mathbb{C} \right\} \cong \mathbb{C}[x]/(x^2).$$

Remark 4.1. These are not arbitrary test cases: $\mathbb{M}_2$ is the smallest semisimple algebra that is noncommutative, while $\mathbf{T}_2$ is the smallest commutative, unital algebra that is not semisimple. (In the context of commutative algebra and deformation theory, it is also known as the algebra of dual numbers over $\mathbb{C}$, since it formalizes the notion of an infinitesimal element vanishing to 2nd order.) Additional motivation comes from results of R. A. J. Howey, who showed that $C^k[0,1]$ is AMNM ($k \geq 1$) ([6]), while observing – in slightly different notation – that the pair $(C^k[0,1], \mathbf{T}_2)$ is not AMNM ([5, Corollary 4.2.5]; see also Remark 4.3 below).

It does not matter which norm we put on $\mathbf{T}_2$ or $\mathbb{M}_2$. To be definite, we give $\mathbb{M}_2$ its natural norm (the C*-algebra norm), but equip $\mathbf{T}_2$ with the norm

$$\left\| \begin{pmatrix} a & b \\ 0 & a \end{pmatrix} \right\|_{\mathbf{T}_2} = |a| + |b|.$$

Theorem 4.2. *Let ω be an unbounded weight function on $\mathbb{N}_{\min}$. Then neither $(\ell^1_\omega(\mathbb{N}_{\min}), \mathbf{T}_2)$ nor $(\ell^1_\omega(\mathbb{N}_{\min}), \mathbb{M}_2)$ are AMNM pairs.*

Proof. Throughout this proof, let A denote the algebra $\ell^1_\omega(\mathbb{N}_{\min})$. For each $m \in \mathbb{N}$ let χ_m be the character

$$\chi_m(k) = \begin{cases} 1 & \text{if } k \geq m, \\ 0 & \text{if } k < m, \end{cases}$$

and let δ_m be the Dirac point mass at m. Set

$$\theta_m = \begin{pmatrix} \chi_m & \omega(m)\delta_m \\ 0 & \chi_m \end{pmatrix} : \mathbb{N} \to \mathbf{T}_2,$$

noting that the range of θ_m consists of commuting elements.

For $1 \leq j \leq k$, we have

$$\begin{aligned} \theta_m(j)\theta_m(k) - \theta_m(j) &= \begin{pmatrix} 0 & \chi_m(j)\omega(m)\delta_m(k) + \omega(m)\delta_m(j)(\chi_m(k) - 1) \\ 0 & 0 \end{pmatrix} \\ &= \begin{pmatrix} 0 & \chi_m(j)\omega(m)\delta_m(k) \\ 0 & 0 \end{pmatrix}. \end{aligned}$$

Hence by symmetry, for general $j, k \in \mathbb{N}_{\min}$ we have

$$\theta_m(j)\theta_m(k) - \theta_m(j \wedge k) = \begin{pmatrix} 0 & \chi_m(j \wedge k)\omega(m)\delta_m(j \vee k) \\ 0 & 0 \end{pmatrix}$$

which vanishes unless $j = k = m$, in which case

$$\theta_m(m)\theta_m(m) - \theta_m(m) = \begin{pmatrix} 0 & \omega(m) \\ 0 & 0 \end{pmatrix}.$$

Thus

$$\operatorname{def}(\theta_m) = \sup_{j,k\in\mathbb{N}} \frac{\|\theta(j)\theta(k) - \theta(j \wedge k)\|}{\omega(j)\omega(k)} = \frac{\omega(m)}{\omega(m)\omega(m)} = \omega(m)^{-1}. \tag{4.1}$$

We now consider the cases of $\mathbf{T}_2$ and $\mathbb{M}_2$ separately. Suppose $\Phi : \mathbb{N}_{\min} \to \mathbf{T}_2$ is multiplicative. Since $\mathbf{T}_2$ has no non-trivial idempotents, this forces $\phi(n)_{12} = 0$ for all n. Hence

$$\|\Phi(m) - \theta_m(m)\| \geq |\Phi(m)_{12} - \theta_m(m)_{12}| \geq \omega(m),$$

so that

$$\operatorname{dist}_{\mathcal{L}(A,\mathbf{T}_2)}(\theta_m, \operatorname{Mult}(A, \mathbf{T}_2)) \geq \sup_{x\in\mathbb{N}} \frac{\|\Phi(x) - \theta_m(x)\|}{\omega(x)} \geq 1 \quad \text{for all } m.$$

Since $\liminf_n \omega(n)^{-1} = 0$, it follows that $(A, \mathbf{T}_2)$ is not an AMNM pair.

A little more work is needed in the case of $\mathbb{M}_2$. Let $C \geq 1 + \omega(1)$. Since $\operatorname{def}(\theta_m) = \omega(m)^{-1}$ is not bounded away from zero, it suffices to prove the following claim.

CLAIM: $\operatorname{dist}_{\mathcal{L}(A,\mathbb{M}_2)}(\theta_m, \operatorname{Mult}(A, \mathbb{M}_2)) \geq C^{-1}$.

Suppose the claim is false. Then there exists $\phi : \mathbb{N}_{\min} \to \mathbb{M}_2$ which is multiplicative (in particular, each $\phi(n)$ is an idempotent in $\mathbb{M}_2$) and satisfies

$$\sup_n \frac{\|\theta_m(n) - \phi(n)\|}{\omega(n)} < C^{-1}. \tag{4.2}$$

Since $\theta_m(1) = I$ and $\|\theta_m(1) - \phi(1)\| \leq C^{-1}\omega(1) < 1$, $\phi(1)$ is an invertible idempotent in $\mathbb{M}_2$, so $\phi(1) = I$. Then, since ϕ is multiplicative, $I = \phi(1) = \phi(1)\phi(m) = \phi(m)$. But then taking $n = m$ in (4.2) and recalling that $\theta_m(m)_{12} = \omega(m)$, we get $1 \leq C^{-1}$, a contradiction. This proves the claim, and hence proves the theorem. □

Remark 4.3. A (bounded) multiplicative function Ψ from a Banach algebra A to $\mathbf{T}_2$ is easily seen to be of the form

$$\Psi(a) = \begin{pmatrix} \varphi(a) & D(a) \\ 0 & \varphi(a) \end{pmatrix},$$

where $\varphi \in \operatorname{Mult}(A, \mathbb{C})$ and $D : A \to \mathbb{C}$ is a (bounded) derivation with respect to the bimodule action of A on $\mathbb{C}$ via ϕ. In the case where A has a dense subspace consisting of commuting idempotents, the only such bounded derivation is 0; and the failure of $(\ell^1_\omega(\mathbb{N}_{\min}), \mathbf{T}_2)$ to be AMNM can be interpreted as saying that there are "approximate point derivations" of large norm on $\ell^1_\omega(\mathbb{N}_{\min})$. This perspective (motivated by cohomological questions about the "Feinstein algebras") was the original approach used to construct the counter-example seen in proving Theorem 4.2. With hindsight, a similar idea can be seen behind Howey's result that $(C^k[0,1], \mathbf{T}_2)$ is not an AMNM pair.

Remark 4.4. If ω is any weight on $\mathbb{N}_{\min}$ with $\liminf_n \omega(n) < \infty$, then $\ell^1_\omega(\mathbb{N}_{\min})$ is known to be approximately amenable (in fact, it is boundedly approximately contractible, by combining the discussion in Example 2.5 with [3, Corollary 4.5]). Thus, Theorem 4.2 shows that we can have approximately amenable Banach algebras A and finite-dimensional Banach algebras B such that (A, B) is not an AMNM pair, in contrast with what happens when A is amenable [10, Theorem 3.1].

Theorem 4.2 suggests that if we seek positive results, we are better off considering the unweighted convolution algebras of semilattices. This is corroborated by the next result.

Theorem 4.5. *Let S be a semilattice. Then $(\ell^1(S), \mathbf{T}_2)$ is an AMNM pair.*

Proof. Write $\theta(e) = \begin{pmatrix} a(e) & b(e) \\ 0 & a(e) \end{pmatrix}$ for functions $a, b : S \to \mathbb{C}$. Since

$$\begin{pmatrix} a(e) & b(e) \\ 0 & a(e) \end{pmatrix} \begin{pmatrix} a(f) & b(f) \\ 0 & a(f) \end{pmatrix} = \begin{pmatrix} a(e)a(f) & a(e)b(f) + b(e)a(f) \\ 0 & a(e)a(f) \end{pmatrix} \tag{4.3}$$

we see that $\operatorname{def}(a) \leq \operatorname{def}(\theta) < 2/9$. By Proposition 3.3, there exists a multiplicative function $\chi : S \to \mathbb{C}$ such that

$$|a(e) - \chi(e)| < 3|a(e)^2 - a(e)| < \frac{3\operatorname{def}(\theta)}{2} < 1/3 \quad \text{for all } e \in S.$$

Note that

$$\begin{aligned} \operatorname{def}(\theta) &= \sup_{e \in S} \|\theta(e)^2 - \theta(e)\| \\ &= \sup_{e \in S} \left\| \begin{pmatrix} a(e)^2 - a(e) & 2a(e) - 1)b(e) \\ 0 & a(e)^2 - a(e) \end{pmatrix} \right\| \\ &= \sup_{e \in S} |a(e)^2 - a(e)| + |2a(e) - 1|\,|b(e)|. \end{aligned}$$

Let $e \in S$. Since $|a(e)^2 - a(e)| \leq \operatorname{def}(a) \leq \operatorname{def}(\theta) < 2/9$, applying Lemma 3.2 gives $\operatorname{dist}_{\mathbb{C}}(a(e), \{0,1\}) < 2/3$, and hence $|2a(e) - 1| > 1/3$. Therefore

$$\|\theta(e) - \chi(e)I\| = |a(e) - \chi(e)| + |b(e)| \leq \frac{3}{2}|a(e)^2 - a(e)| + 3|(2a(e) - 1)b(e)| \leq 3\operatorname{def}(\theta).$$

□

5 $(\ell^1(S), \mathbb{M}_2)$ IS AMNM FOR ANY SEMILATTICE S

For reasons of technical convenience, we shall work mostly with the Hilbert-Schmidt norm on $\mathbb{M}_2$, defined by $\|A\|_{\mathrm{HS}}^2 = \operatorname{Tr}(A^*A)^{1/2}$. It might be conceptually clearer to use the operator norm throughout, but this seems to yield worse constants in later inequalities, which are obtained by bootstrapping up the earlier ones.

Notation. If $A \in \mathbb{M}_2$ and $\varepsilon > 0$ let

$$\overline{\mathbb{B}}_A^{\mathrm{HS}}(\varepsilon) = \{B \in \mathbb{M}_2 \ : \ \|A - B\|_{\mathrm{HS}} \leq \varepsilon\}.$$

(the closed ball of radius ε centreed on A, in the Hilbert-Schmidt norm).

Theorem 5.1. *Let $\delta < 1/36$, and let $\theta : S \to \mathbb{M}_2$ be a function satisfying*

$$\sup_{e.f\in S} \|\theta(e)\theta(f) - \theta(ef)\|_{\mathrm{HS}} \leq \delta. \tag{5.1}$$

Then there exists a (bounded) multiplicative function $\phi : S \to \mathbb{M}_2$ such that

$$\sup_{x\in S} \|\theta(x) - \phi(x)\|_{\mathrm{HS}} \leq 18\delta.$$

In particular, $(\ell^1(S), \mathbb{M}_2)$ is a strongly AMNM pair.

Note that we do not assume in (5.1) that $\sup_{e\in S} \|\theta(e)\|_{\mathrm{HS}} < \infty$, but that this will emerge during the proof (see Proposition 5.4).

5.1 MOTIVATING THE PROOF OF THEOREM 5.1

Let S be a semilattice. Our proof that $(\ell^1(S), \mathbb{C})$ is a uniformly AMNM pair can be broken down into three steps:

1. Show that there is a constant c, such that whenever $\theta : S \to \mathbb{C}$ is δ-multiplicative, $\theta(S) \subseteq \overline{\mathbb{D}}_0(c\delta) \cup \overline{\mathbb{D}}_1(c\delta)$. (In effect, this step "approximately discretizes" the problem.)
2. Put $S_k := \theta^{-1}(\overline{\mathbb{D}}_k(c\delta))$ for $k = 0, 1$, so that S is partitioned as $S_1 \cup S_0$. Show that $S_1 \cdot S_1 \subseteq S_1$, $S_1 \cdot S_0 \subseteq S_0$, and $S_0 \cdot S_0 \subseteq S_0$. (Although we did not do these calculations explicitly, they are implicit in the process of checking S_1 is a filter in S.)
3. Define $\phi : S \to \{0, 1\}$ by $\phi = 1$ on S_1 and $\phi = 0$ on S_0. By the first step, $\|\phi - \theta\|_\infty \leq c\delta$, and by the second step, ϕ is multiplicative.

The strategy we shall adopt is to mimic each of these steps, but now allow our maps to take values in $\mathbb{M}_2$ rather than $\mathbb{C}$. As a first step, we need some characterization of multiplicative functions $\phi : S \to \mathbb{M}_2$, which reduces down to the problem of describing the possible semilattices inside the multiplicative semigroup $\mathbb{M}_2$. This is not too hard, once we make the following observation: *if $P \in \mathbb{M}_2$ is a rank-one idempotent, then the only idempotents which commute with P are I, P, $I - P$ and* 0. (We will see later that there is an "approximate version" of this.)

Secondly, observe that if θ is – as claimed – a perturbation of a multiplicative function ϕ, then by the previous remarks $\theta(S)$ should be contained in a small-ball neighbourhood of $\phi(S)$, which in turn is contained in a set of at most four commuting idempotents. To prove Theorem 5.1, we reverse this line of reasoning, and identify a commuting set L of idempotents in $\mathbb{M}_2$, a small-ball neighbourhood of which will contain $\theta(S)$. (See Proposition 5.4 for the details.) Then, since the elements of L are well-separated, there is only one realistic candidate for the map ϕ: namely, it should send a given $x \in S$ to the element of L nearest to $\theta(x)$. This map ϕ will clearly be close in norm to θ, so all that will remain is to check that ϕ is multiplicative: this can be done through a case-by-case analysis, although some work is needed since we do not assume $\sup_{e\in S} \|\theta(e)\|_{\mathrm{HS}} < \infty$.

To identify the set L, we make heavy use of a small but technical result, based on the following idea: the trace of an approximately idempotent 2×2 matrix must be close to an integer, which then equalse the rank of any nearby idempotent. This will be made precise in the next lemma.

5.2 A KEY TECHNICAL LEMMA

The following lemma is our basic tool for working with approximately idempotent elements of $\mathbb{M}_2$. It is here that the Hilbert-Schmidt norm seems to be convenient.

Lemma 5.2 (Key estimates)**.** *Let* $A \in \mathbb{M}_2$*. Suppose* $\|A - A^2\|_{\mathrm{HS}} \leq \varepsilon < 2/9$*. Then* $\|2A - I\|_{\mathrm{HS}} \geq \sqrt{2 - 6\varepsilon}$*. Moreover, exactly one of the following three possibilities occurs:*

(i) $\mathrm{Tr}(A) \in \overline{\mathbb{D}}_2(3\varepsilon/\sqrt{2})$, *and* $A \in \overline{\mathbb{B}}_I^{\mathrm{HS}}(2\varepsilon)$;

(ii) $\mathrm{Tr}(A) \in \overline{\mathbb{D}}_1(2\varepsilon)$, *and* $A \in \overline{\mathbb{B}}_P^{\mathrm{HS}}(3\varepsilon/2)$ *for some rank-*1 *idempotent* $P \in \mathbb{M}_2$;

(iii) $\mathrm{Tr}(A) \in \overline{\mathbb{D}}_0(3\varepsilon/\sqrt{2})$, *and* $A \in \overline{\mathbb{B}}_0^{\mathrm{HS}}(2\varepsilon)$.

In case (i)*,* A *is invertible and* $\|A^{-1}\|_{\mathrm{HS}} \leq 3$*. In case* (iii)*,* $I - A$ *is invertible and* $\|(I - A)^{-1}\|_{\mathrm{HS}} \leq 3$*.*

Proof. By conjugating with an appropriate unitary matrix, we may assume without loss of generality that A is upper triangular, say $A = \begin{pmatrix} a & b \\ 0 & d \end{pmatrix}$. Then

$$A - A^2 = \begin{pmatrix} a - a^2 & b(1 - a - d) \\ 0 & d - d^2 \end{pmatrix}$$

so that $|a - a^2|^2 + |b(1 - a - d)|^2 + |d - d^2|^2 \leq \|A - A^2\|_{\mathrm{HS}}^2 \leq \varepsilon^2$. In particular, by the scalar case (Lemma 3.2),

$$\mathrm{dist}_{\mathbb{C}}(a, \{0,1\}) \leq \frac{3}{2}|a - a^2| \quad \text{and} \quad \mathrm{dist}_{\mathbb{C}}(d, \{0,1\}) \leq \frac{3}{2}|d - d^2|,$$

implying that

$$\mathrm{dist}_{\mathbb{C}}(a + d, \{0,1,2\}) \leq \frac{3}{2}|a - a^2| + \frac{3}{2}|d - d^2| \leq \frac{3}{\sqrt{2}}\left(|a - a^2|^2 + |d - d^2|^2\right)^{1/2} \leq \frac{3}{\sqrt{2}}\varepsilon.$$

Moreover,

$$\begin{aligned} \|2A - I\|_{\mathrm{HS}}^2 &\geq |2a - 1|^2 + |2d - 1|^2 \\ &\geq 2 - 4|a - a^2| - 4|d - d^2| \\ &\geq 2 - 4\sqrt{2}\left(|a - a^2|^2 + |d - d^2|^2\right)^{1/2} \quad \geq 2 - 4\sqrt{2}\varepsilon \geq 2 - 6\varepsilon. \end{aligned}$$

Note that since $\varepsilon < 2/9$, we have

$$\frac{3\varepsilon}{\sqrt{2}} < \frac{\sqrt{2}}{3} \quad \text{and} \quad \frac{3\varepsilon}{2} < \frac{1}{3}.$$

THE CASE $|a+d-2| \le 2\varepsilon$. We have

$$|1-a-d| \ge 1 - 3\varepsilon/\sqrt{2} > \frac{1}{2},$$

and so $|b| \le 2|b(1-a-d)|$. We also have $|a-1| \le \frac{3}{2}|a-a^2|$ and $|d-1| \le \frac{3}{2}|d-d^2|$. Therefore

$$\begin{aligned}\|A-I\|_{\mathrm{op}} \le \|A-I\|_{\mathrm{HS}} &= \left(|a-1|^2+|b|^2+|d-1|^2\right)^{1/2} \\ &\le 2\left(|a-a^2|^2+|b(1-a-d)|^2+|d-d^2|^2\right)^{1/2} \le 2\varepsilon < \frac{4}{9}.\end{aligned}$$

This gives

$$\|A^{-1}\|_{\mathrm{HS}} \le \sqrt{2}\|A^{-1}\|_{\mathrm{op}} \le \frac{\sqrt{2}\cdot 9}{5} < 3.$$

THE CASE $|a+d-1| \le 2\varepsilon$ There are two sub-cases: either

(a) $|a-1| \le \frac{3}{2}|a^2-a|$ and $|d| \le \frac{3}{2}|d^2-d|$; or

(b) $|a| \le \frac{3}{2}|a^2-a|$ and $|d-1| \le \frac{3}{2}|d^2-d|$.

Suppose we are in case (a). Put $P = \begin{pmatrix} 1 & b \\ 0 & 0 \end{pmatrix}$, which is clearly an idempotent; then

$$\|A-P\|_{\mathrm{HS}}^2 = |a-1|^2+|d|^2 \le \frac{9}{4}|a^2-a|^2 + \frac{9}{4}|d^2-d|^2 \le \frac{9}{4}\|A\|_{\mathrm{HS}}^2$$

as claimed. If we are in case (b), put $P = \begin{pmatrix} 0 & b \\ 0 & 1 \end{pmatrix}$; a similar calculation then shows that $\|A-P\|_{\mathrm{HS}} \le \frac{3}{2}\varepsilon$ as before.

THE CASE $|a+d| \le 2\varepsilon$ In this case, we also have $|1-a-d| > 1/2$, so

$$|b| \le 2|b(1-a-d)|.$$

We also have

$$|a| \le \frac{3}{2}|a-a^2| \quad \text{and} \quad |d| \le \frac{3}{2}|d-d^2|,$$

so that a similar estimate to the first case yields $\|A\|_{\mathrm{HS}} \le 2\varepsilon$. ☐

5.3 THE PROOF OF THEOREM 5.1

To save needless repetition, we will assume for the rest of this section that $0 \le \delta < 1/36$.

Let $\theta : S \to \mathbb{M}_2$ satisfy the condition (5.1). To simplify formulas, we shall use the following abbreviations: $\widetilde{e}$ stands for $\theta(e)$, $\widetilde{f}$ for $\theta(f)$, $\widetilde{ef}$ for $\theta(ef)$, and so forth.

We start by taking $e = f$ in condition (5.1) and using Lemma 5.2 with $\varepsilon = \delta$. This gives

$$\operatorname{Tr}\theta(S) \subseteq \overline{\mathbb{D}}_0(3\delta/\sqrt{2}) \cup \overline{\mathbb{D}}_1(3\delta/\sqrt{2}) \cup \overline{\mathbb{D}}_2(3\delta/\sqrt{2}). \tag{5.2}$$



Figure 1: Dependencies between results in this section. (P = Proposition; L = Lemma.)

Define $S_k := (\mathrm{Tr}\circ\theta)^{-1}(\overline{\mathbb{D}}_k(3\delta/\sqrt{2}))$ for $k = 0, 1, 2$. Since $3\delta/\sqrt{2} < 1/16$, (5.2) implies that $\mathrm{Tr}\,\theta(S_k) \subseteq \mathbb{D}_k(1/16)$ for $k \in \{0, 1, 2\}$. In particular,

$$\text{if } g \in S, k \in \mathbb{Z} \text{ and } |\mathrm{Tr}\,\theta(g) - k| < 0.9, \text{ then } g \in S_k. \tag{5.3}$$

By cases (i) and (iii) of Lemma 5.2,

$$S_2 = \theta^{-1}(\overline{\mathbb{B}}_I^{\mathrm{HS}}(2\delta)) \quad \text{and} \quad S_0 = \theta^{-1}(\overline{\mathbb{B}}_0^{\mathrm{HS}}(2\delta)).$$

Define $\phi : S_2 \sqcup S_0 \to \mathbb{M}_2$ by $\phi(S_0) = \{0\}$ and $\phi(S_2) = \{I\}$. Then by construction,

$$\sup_{x \in S_2 \sqcup S_0} \|\widetilde{x} - \phi(x)\|_{\mathrm{HS}} \le 2\delta. \tag{5.4}$$

Proposition 5.3 (Some easy properties)**.** *Let $e, f \in S$.*

(i) *If $e, f \in S_2$ then $ef \in S_2$.*

(ii) *If $e \in S_0$ and $f \in S$ then $ef \in S_0$.*

In particular, $S_2 \sqcup S_0$ is a subsemigroup of S, and $\phi : S_2 \sqcup S_0 \to \mathbb{M}_2$ is multiplicative.

Proof. Let $e, f \in S$. Then

$$\|(\widetilde{ef} - \widetilde{e})\widetilde{f}\|_{\mathrm{HS}} \le \|\widetilde{ef}\widetilde{f} - ef\|_{\mathrm{HS}} + \|\widetilde{ef} - \widetilde{e}\widetilde{f}\|_{\mathrm{HS}} \le 2\delta.$$

If $f \in S_2$, then by Lemma 5.2(i) with $\varepsilon = \delta$ we have $\|f^{-1}\|_{\mathrm{HS}} \le 3$. Hence by Cauchy-Schwarz

$$|\mathrm{Tr}(\widetilde{ef} - \widetilde{e})| \le \|(\widetilde{ef} - \widetilde{e})\widetilde{f}\|_{\mathrm{HS}}\, \|\widetilde{f}^{-1}\|_{\mathrm{HS}} \le 6\delta. \tag{5.5}$$

Therefore, if $e \in S_2$,

$$|\mathrm{Tr}(\widetilde{ef}) - 2| \leq \frac{20\delta}{9} + 6\delta < \frac{5}{81} + \frac{1}{6} < \frac{11}{48}.$$

So by the observation (5.3), $ef \in S_2$, and (i) is proved.

Also, note that $\|\widetilde{e}\,\widetilde{ef} - \widetilde{ef}\|_{\mathrm{HS}} \leq \delta$. If $e \in S_0$, $\|\widetilde{e} - I\|_{\mathrm{HS}} \leq 3$ by Lemma 5.2, and so by Cauchy-Schwarz,

$$|\mathrm{Tr}\,\widetilde{ef}| \leq \|(\widetilde{e} - I)^{-1}\|_{\mathrm{HS}}\,\|(\widetilde{e} - I)\widetilde{ef}\|_{\mathrm{HS}} \leq 3\delta < \frac{1}{12}.$$

Hence $ef \in S_0$. This proves (ii). The final statement of the proposition now follows easily from (i) and (ii). $\square$

Combining (5.4) and Proposition 5.3, we get a proof of Theorem 5.1 in the special case where $S_1 = \emptyset$. *We shall therefore assume, for the rest of this section, that* S_1 *is non-empty.*

Proposition 5.4. *There exists a rank* 1*-idempotent* $P \in \mathbb{M}_2$ *such that*

$$\theta(S_1) \subseteq \overline{\mathbb{B}}^{\mathrm{HS}}_P(18\delta) \sqcup \overline{\mathbb{B}}^{\mathrm{HS}}_{I-P}(18\delta).$$

Moreover, given $e, f \in S_1$*:*

- *if* $\widetilde{e}, \widetilde{f} \in \overline{\mathbb{B}}^{\mathrm{HS}}_P(18\delta)$*, then so does* $\widetilde{ef}$*;*
- *if* $\widetilde{e}, \widetilde{f} \in \overline{\mathbb{B}}^{\mathrm{HS}}_{I-P}(18\delta)$*, then so does* $\widetilde{ef}$*;*
- *if* $\widetilde{e} \in \overline{\mathbb{B}}^{\mathrm{HS}}_P(18\delta)$ *and* $\widetilde{f} \in \overline{\mathbb{B}}^{\mathrm{HS}}_{I-P}(18\delta)$*, then* $ef \in S_0$*.*

The proof of Proposition 5.4 requires some work, which we break up into several lemmas. The first of these is an "approximate version" of the following observation: if P and Q are rank-one idempotents in $\mathbb{M}_2$ with $PQ = 0 = QP$, then $P + Q = I$. It also shows where we first use the condition that $\delta < 1/36$, as this is needed in order to bootstrap up from Lemma 5.2.

Lemma 5.5. *Let* $e, f \in S_1$ *with* $ef \in S_0$*. Then* $\widetilde{e} + \widetilde{f} \in \overline{\mathbb{B}}^{\mathrm{HS}}_I(16\delta)$*, and* $\|\widetilde{e} - \widetilde{f}\|_{\mathrm{HS}} \geq \sqrt{11/6} - 16\delta$*.*

Proof. Let $A = \widetilde{e} + \widetilde{f}$. We wish to prove A is close to I; this will follow if we can show A is approximately idempotent and has trace close to 2.

Since $ef \in S_0$, $\|\widetilde{ef}\|_{\mathrm{HS}} \leq 2\delta$ (by applying Lemma 5.2 with $\varepsilon = \delta$). By (5.1), this gives

$$\|\widetilde{e}\widetilde{f} + \widetilde{f}\widetilde{e}\|_{\mathrm{HS}} \leq 2\delta + 2\|\widetilde{ef}\|_{\mathrm{HS}} \leq 6\delta.$$

Hence

$$\begin{aligned} \|A^2 - A\|_{\mathrm{HS}} &= \|\widetilde{e} + \widetilde{f} - (\widetilde{e} + \widetilde{f})^2\|_{\mathrm{HS}} \\ &\leq \|\widetilde{e}^2 - \widetilde{e}\|_{\mathrm{HS}} + \|\widetilde{f}^2 - \widetilde{f}\|_{\mathrm{HS}} + \|\widetilde{e}\widetilde{f} + \widetilde{f}\widetilde{e}\|_{\mathrm{HS}} \leq 8\delta \leq \frac{2}{9}. \end{aligned}$$

By Lemma 5.2 with $\varepsilon = 8\delta$, we therefore have

$$\mathrm{Tr}(A) \in \bigcup_{k=0}^{2} \overline{\mathbb{D}}_k(160\delta/9) \subseteq \bigcup_{k=0}^{2} \overline{\mathbb{D}}_k(40/81).$$

On the other hand, since $\operatorname{Tr}\widetilde{e}, \operatorname{Tr}\widetilde{f} \in \overline{\mathbb{D}}_1(20\delta/9)$, we have

$$\operatorname{Tr}(A) \in \overline{\mathbb{D}}_2(40\delta/9) \subseteq \overline{\mathbb{D}}_2(10/81).$$

So we must be in case (i) of Lemma 5.2, which yields $\|A - I\|_{\mathrm{HS}} \leq 2\varepsilon = 16\delta$ as claimed. Finally, since $e \in S_1$,

$$\|2\widetilde{e} - I\|_{\mathrm{HS}} \geq \sqrt{2 - 6\delta} \geq \sqrt{11/6}\,,$$

and so

$$\|\widetilde{e} - \widetilde{f}\|_{\mathrm{HS}} \geq \|2\widetilde{e} - I\|_{\mathrm{HS}} - \|\widetilde{e} + \widetilde{f} - I\|_{\mathrm{HS}} > \sqrt{11/6} - 16\delta.$$

☐

Intuitively, we should have $S_1 \cdot S_1 \subseteq (S_1 \cup S_0)$ since elements of $\theta(S_k)$ are close to idempotents of rank k. Some care is needed to show this, because we have no *a priori* upper bound on norms of elements in $\theta(S_1)$.

Lemma 5.6.

(i) *If $e \in S$, $f \in S_2$ and $e \succeq f$, then $e \in S_2$.*

(ii) $S_1 \cdot S_1 \subseteq S_1 \cup S_0$.

Proof. Let $e, f \in S$ with $ef = f$. Then $\|\widetilde{e}\widetilde{f} - \widetilde{f}\|_{\mathrm{HS}} \leq \delta$. If $f \in S_2$, then $\|\widetilde{f}^{-1}\|_{\mathrm{HS}} \leq 3$ by Lemma 5.2. Hence, by Cauchy-Schwarz,

$$|\operatorname{Tr}(\widetilde{e} - I)| \leq \|\widetilde{e}\widetilde{f} - \widetilde{f}\|_{\mathrm{HS}} \|\widetilde{f}^{-1}\|_{\mathrm{HS}} \leq 3\delta < \frac{1}{12}$$

so that $e \in S_2$. This proves (i). Now if $e, g \in S_1$, put $f = eg$; by part (i) we cannot have $f \in S_2$, so (ii) is proved. ☐

To analyze S_1 in further detail, the following lemma is useful.

Lemma 5.7 (Chains in S_1)**.** *Let $e, f \in S_1$ with $e \succeq f$. Then $\|\widetilde{e} - \widetilde{f}\|_{\mathrm{HS}} \leq 8\delta$.*

Proof. Put $A = \widetilde{e} - \widetilde{f}$. We show that A is approximately idempotent and has small trace, so must be close to 0 by Lemma 5.2. In detail: observe that

$$\|A^2 - A\|_{\mathrm{HS}} = \|\widetilde{e}^2 - \widetilde{e} + \widetilde{f}^2 - \widetilde{f} - \widetilde{e}\widetilde{f} + \widetilde{f} - \widetilde{f}\widetilde{e} + \widetilde{f}\|_{\mathrm{HS}} \leq 4\delta$$

and $\operatorname{Tr}(A) \leq |1 - \operatorname{Tr}\widetilde{e}| + |1 - \operatorname{Tr}\widetilde{f}| \leq 4\delta$. Using case (iii) of Lemma 5.2, with $\varepsilon = 4\delta < 1/9$, we get $\|\widetilde{e} - \widetilde{f}\|_{\mathrm{HS}} \leq 2\varepsilon = 8\delta$. ☐

Proof of Proposition 5.4. Recall that S_1 is, by assumption, non-empty. Consider the relation on S_1 defined by $\{(e, f) \in S_1 \times S_1 \;:\; ef \in S_1\}$, and denote it by $\sim$. Clearly $\sim$ is symmetric and reflexive. If $e \sim f$, then $ef \in S_1$ with $e \succeq ef$ and $f \succeq ef$, so applying Lemma 5.7 we have

$$\|\widetilde{e} - \widetilde{f}\|_{\mathrm{HS}} \leq \|\widetilde{e} - \widetilde{ef}\|_{\mathrm{HS}} + \|\widetilde{f} - \widetilde{ef}\|_{\mathrm{HS}} \leq 16\delta. \tag{5.6}$$

Suppose $e \sim f$ and $f \sim g$. Then, since $\frac{4}{3} > 48\delta$, we have

$$\|\widetilde{e} - \widetilde{g}\|_{\mathrm{HS}} \leq \|\widetilde{e} - \widetilde{f}\|_{\mathrm{HS}} + \|\widetilde{f} - \widetilde{g}\|_{\mathrm{HS}} \leq 32\delta < \frac{4}{3} - 16\delta < \sqrt{11/6} - 16\delta.$$

Hence, by (the contrapositive of) Lemma 5.5, $eg \notin S_0$. Therefore, by Lemma 5.6, $eg \in S_1$, i.e. $e \sim g$. This shows that $\sim$ is transitive, and hence it is an equivalence relation on S_1. A little thought shows that each equivalence class for this relation is closed under multiplication.

Now let $e, f, g \in S_1$. Suppose $e \not\sim f$ and $f \not\sim g$. Then, since $ef \in S_0$ and $fg \in S_0$, Lemma 5.5 gives $\|\widetilde{e} + \widetilde{f} - I\|_{\mathrm{HS}} \leq 16\delta$ and $\|\widetilde{f} + \widetilde{g} - I\|_{\mathrm{HS}} \leq 16\delta$. Thus

$$\|\widetilde{e} - \widetilde{g}\|_{\mathrm{HS}} \leq 32\delta \leq 4/3 - 16\delta;$$

arguing as above, we deduce that $e \sim g$. Thus there are at most two equivalence classes for this relation.

By assumption S_1 is non-empty; fix $p_0 \in S_1$. By Lemma 5.2, there exists a rank-1 idempotent $P \in \mathbb{M}_2$ such that $\widetilde{p}_0 \in \overline{\mathbb{B}}_P^{\mathrm{HS}}\left(\frac{3}{2}\delta\right)$. Let $x \in S_1$. If $x \sim p_0$, then by (5.6) we have

$$\|\widetilde{x} - P\|_{\mathrm{HS}} \leq \|\widetilde{x} - \widetilde{p}_0\|_{\mathrm{HS}} + \|\widetilde{p}_0 - P\|_{\mathrm{HS}} \leq 18\delta.$$

If $x \not\sim p_0$, then since $xp_0 \in S_0$, Lemma 5.5 gives $\|\widetilde{x} - \widetilde{p}_0 + I\|_{\mathrm{HS}} \leq 16\delta$, and so

$$\|\widetilde{x} - (I - P)\|_{\mathrm{HS}} \leq \|\widetilde{x} - \widetilde{p}_0 + I\|_{\mathrm{HS}} + \|\widetilde{p}_0 - P\|_{\mathrm{HS}} \leq 18\delta.$$

Finally, let $e, f \in S_1$.

- If $\widetilde{e}, \widetilde{f} \in \overline{\mathbb{B}}_P^{\mathrm{HS}}(18\delta)$ then $e \sim p_0 \sim f$, so that $ef \sim p_0$ (as equivalence classes are closed under multiplication) and therefore $\widetilde{ef} \in \overline{\mathbb{B}}_{I-P}^{\mathrm{HS}}(18\delta)$.
- If $\widetilde{e}, \widetilde{f} \in \overline{\mathbb{B}}_{I-P}^{\mathrm{HS}}(18\delta)$, then $e \not\sim p_0$, $f \not\sim p_0$; since there are at most two equivalence classes, $e \sim f$. Thus $e \sim ef \sim f$, so $ef \not\sim p_0$, so $\widetilde{ef} \in \overline{\mathbb{B}}_{I-P}^{\mathrm{HS}}(18\delta)$.
- If $\widetilde{e} \in \overline{\mathbb{B}}_P^{\mathrm{HS}}(18\delta)$ and $\widetilde{f} \in \overline{\mathbb{B}}_{I-P}^{\mathrm{HS}}(18\delta)$, then $e \not\sim f$; hence $ef \in S_0$ as claimed.

□

We now *fix* an idempotent $P \in \mathbb{M}_2$ that satisfies the conclusions of Proposition 5.4. The sets

$$\overline{\mathbb{B}}_I^{\mathrm{HS}}(2\delta) \quad , \quad \overline{\mathbb{B}}_P^{\mathrm{HS}}(18\delta) \quad , \quad \overline{\mathbb{B}}_{I-P}^{\mathrm{HS}}(18\delta) \quad , \quad \overline{\mathbb{B}}_0^{\mathrm{HS}}(2\delta)$$

are pairwise disjoint, and their union contains $\theta(S)$. Recall that we have already defined $\phi : S_2 \sqcup S_0 \to \mathbb{M}_2$ which is multiplicative; now define $\phi : S_1 \to \mathbb{M}_2$ by setting $\phi(x)$ to be whichever of P and $I - P$ is closer to $\widetilde{x}$. Explicitly, the map $\phi : S \to \mathbb{M}_2$ satisfies:

$$\phi(x) = \begin{cases} I & \text{if } x \in S_2, \\ P & \text{if } \widetilde{x} \in \overline{\mathbb{B}}_P^{\mathrm{HS}}(18\delta), \\ I - P & \text{if } \widetilde{x} \in \overline{\mathbb{B}}_{I-P}^{\mathrm{HS}}(18\delta), \\ 0 & \text{if } x \in S_0. \end{cases}$$

By construction, $\|\widetilde{x} - \phi(x)\|_{\mathrm{HS}} \leq 18\delta$ for all $x \in S$, and so it remains to show ϕ is multiplicative.

Let $S_p := \theta^{-1}(\overline{\mathbb{B}}_P^{\mathrm{HS}}(18\delta))$ and $S_q := \theta^{-1}(\overline{\mathbb{B}}_{I-P}^{\mathrm{HS}}(18\delta))$; we have $S_1 = S_p \sqcup S_q$. To show that ϕ is multiplicative, it sufffices to verify the following claims:

1. $S_2 \cdot S_2 \subseteq S_2$.
2. $S \cdot S_0 \subseteq S_0$.
3. $S_2 \cdot S_p \subseteq S_p$ and $S_2 \cdot S_q \subseteq S_q$.
4. $S_p \cdot S_p \subseteq S_p$ and $S_q \cdot S_q \subseteq S_q$.
5. $S_p \cdot S_q \subseteq S_0$.

Assertions 1 and 2 follow from Proposition 5.3. Assertions 4 and 5 follow from Proposition 5.4. Assertion 3 requires some more work, and is dealt with in our final proposition.

Proposition 5.8. *Let $e \in S_2$. If $f \in S_p$, then so is ef; if $f \in S_q$, then so is ef.*

Remark 5.9. Although we know that $\widetilde{e}$ is close to I for each $e \in S_2$, there might still exist rank-1 idempotents $R \in \mathbb{M}_2$ such that $\|\widetilde{e} \cdot R - R\|$ is large. So while the conclusion of Proposition 5.8 is as one would expect, the proof is somewhat circuitous.

Proof of Proposition 5.8. Let $e \in S_2$ and $f \in S_1$. By Lemma 5.6, $ef \notin S_2$. We claim that $ef \notin S_0$. For, assume $ef \in S_0$: then by Lemma 5.2, $\|\widetilde{ef}\|_{\mathrm{HS}} \leq 2\delta$, so using Cauchy-Schwarz and Lemma 5.2(i), we get

$$|\mathrm{Tr}\,\widetilde{f}| \leq \|\widetilde{e}^{-1}\|_{\mathrm{HS}} \|\widetilde{e}\widetilde{f}\|_{\mathrm{HS}} \leq 3(\delta + \|\widetilde{ef}\|_{\mathrm{HS}}) \leq 9\delta < \frac{1}{4}.$$

But, by (5.3), this implies $f \in S_0$, contradicting the assumption that $f \in S_1$. The only remaining possibility is that $ef \in S_1$.

Now recall the equivalence relation from the proof of Proposition 5.4. If $x, y \in S_1$ with $x \succeq y$, then $x \sim y$ (by definition of that relation). So if e and f are as above, $f \sim ef$. We also saw in the proof of Proposition 5.4 that if $x, y \in S_1$ with $x \sim y$, then either both belong to S_p (when $x \sim y \sim p_0$) or both belong to S_q (when $x \not\sim p_0$ and $y \not\sim p_0$). This completes the proof. $\square$

Since Proposition 5.8 implies Assertion 3, the function $\phi : S \to \mathbb{M}_2$ is indeed multiplicative, and this completes the proof of Theorem 5.1.

6 CONCLUDING REMARKS AND QUESTIONS

It would be interesting to try and find an intrinsic condition on a semilattice which is necessary and sufficient for the existence of some weight ω such that $\ell^1_\omega(S)$ is not AMNM. We have seen (Example 3.12(ii) and Theorem 3.13) that $\mathrm{prol}(S) = +\infty$ is a necessary condition; it may also be a sufficient condition, although we have not investigated further.

We leave AMNM pair problems for $(\ell^1(S), B)$, with B an arbitrary Banach algebra, for future work. There is some evidence to suggest that the method used for $B = \mathbb{M}_2$ can be extended to $B = \mathbb{M}_n$, $n \geq 3$, although a more laborious case-by-case analysis would be required. (However, see Remark 1.2.) A more streamlined and unified proof would then be desirable. Finally, we close with the following question: *is $(\ell^1(S), \mathcal{B}(E))$ AMNM for every Banach space E? If not, what if we restrict to the cases $E = \ell^p$ for $1 < p < \infty$?*

ACKNOWLEDGEMENTS

Some parts of this paper are based on work done in 2011, during a visit to the University of Newcastle upon Tyne. The author thanks the School of Mathematics and Statistics at the University for their hospitality.

The work here was supported by NSERC Discovery Grant 402153-2011.

REFERENCES

[1] H. G. DALES AND R. J. LOY, *Approximate amenability of semigroup algebras and Segal algebras*, Dissertationes Math. (Rozprawy Mat.), 474 (2010), p. 58.

[2] J. F. FEINSTEIN, *Strong Ditkin algebras without bounded relative units*, Int. J. Math. Math. Sci., 22 (1999), pp. 437–443.

[3] F. GHAHRAMANI, R. J. LOY, AND Y. ZHANG, *Generalized notions of amenability, II.*, J. Funct. Anal., 254 (2008), pp. 1776–1810.

[4] E. HEWITT AND H. S. ZUCKERMAN, *The l_1-algebra of a commutative semigroup*, Trans. Amer. Math. Soc., 83 (1956), pp. 70–97.

[5] R. A. J. HOWEY, *Approximately multiplicative maps between some Banach algebras*, PhD thesis, University of Newcastle upon Tyne, 2000.

[6] ——, *Approximately multiplicative functionals on algebras of smooth functions*, J. London Math. Soc. (2), 68 (2003), pp. 739–752.

[7] K. JAROSZ, *Perturbations of Banach algebras*, vol. 1120 of Lecture Notes in Mathematics, Springer-Verlag, Berlin, 1985.

[8] ——, *Almost multiplicative functionals*, Studia Math., 124 (1997), pp. 37–58.

[9] B. E. JOHNSON, *Approximately multiplicative functionals*, J. London Math. Soc. (2), 34 (1986), pp. 489–510.

[10] ——, *Approximately multiplicative maps between Banach algebras*, J. London Math. Soc. (2), 37 (1988), pp. 294–316.

[11] J. LAWRENCE, *The stability of multiplicative semigroup homomorphisms to real normed algebras. I*, Aequationes Math., 28 (1985), pp. 94–101.

[12] S. J. SIDNEY, *Are all uniform algebras AMNM?*, Bull. London Math. Soc., 29 (1997), pp. 327–330.